

Protocol:	The student will: <ul style="list-style-type: none"> Set-up the equipment and supplies, procedure will be simulated on a manikin. 					
Time Permitted:	<table border="1"> <tr> <td>Set-up</td> <td>10 minutes</td> </tr> <tr> <td>Practice Time</td> <td>15 minutes</td> </tr> </table>	Set-up	10 minutes	Practice Time	15 minutes	
Set-up	10 minutes					
Practice Time	15 minutes					
Equipment and Supplies:	See Equipment and Supplies Binder					

Periodontal Dressing (Application & Removal):		
1.1 Assembled and set-up all required: Dental Materials, Instruments and Equipment before the start of procedure.		
1.2 Explained procedure to patient-in lay terms, (communicate throughout appointment) (Say something like: we're placing a dressing to protect the surgical site)		
PREPARE PRODUCT		
1.3 Dispensed product according to MIFU (If applicable).		
1.4 Mixed product according to MIFU (If applicable).		
1.5 Rolled into strips approximately the length of surgical site.		
PLACEMENT		
1.6 Placed small triangle-shaped pieces into the interproximal spaces of the surgical site.		
1.7 Adapted one end of the strip around distal surface of the last tooth in the surgical site.		
1.8 Brought the remainder of the strip forward along facial surface, and gently pressed along the incised gingival margin.		
1.9 Gently pressed the strip into the interproximal areas.		
1.10 Applied the second strip in the same manner from the lingual side.		
1.11 Joined the facial and lingual strips at the distal surface of the last tooth at both ends of the surgical site.		
1.12 Checked for overextension of dressing and interference with occlusion. Removed excess if required. Adjusted the margin to ensure dressing is not rough.		
1.13 Provided post operative instructions: <ul style="list-style-type: none"> dressing will harden in about 2 hours, avoid disturbing, 		

Periodontal Dressing (Application & Removal):	
<ul style="list-style-type: none">• OK if small parts chip off, but contact office if large portion comes off,• OHI: Brush top of dressing lightly with a soft toothbrush.	

REMOVAL (following appropriated time for healing)	
1.14	Checked chart to determine presence of sutures.
1.15	Gently inserted appropriate instrument under the margin.
1.16	Used lateral pressure to pry dressing away from tissue.
1.17	If sutures were present, gently cut the sutures and removed from tissue.
1.18	Gently flossed interproximal to remove any fragments of dressing.
1.19	Irrigated area with warm saline solution to remove superficial debris (used saliva ejector or HVE to remove fluid from mouth).

